

SLC22 family of organic cation and anion transporters (version 2019.4) in the IUPHAR/BPS Guide to Pharmacology Database

Bruno Hagenbuch¹

1. University of Kansas, USA

Abstract

The SLC22 family of transporters is mostly composed of non-selective transporters, which are expressed highly in liver, kidney and intestine, playing a major role in drug disposition. The family may be divided into three subfamilies based on the nature of the substrate transported: organic cations (OCTs), organic anions (OATs) and organic zwitterion/cations (OCTN). Membrane topology is predicted to contain 12 TM domains with intracellular termini, and an extended extracellular loop at TM 1/2.

Contents

This is a citation summary for SLC22 family of organic cation and anion transporters in the [Guide to Pharmacology](#) database (GtoPdb). It exists purely as an adjunct to the database to facilitate the recognition of citations to and from the database by citation analyzers. Readers will almost certainly want to visit the relevant sections of the database which are given here under database links.

[GtoPdb](#) is an expert-driven guide to pharmacological targets and the substances that act on them. GtoPdb is a reference work which is most usefully represented as an on-line database. As in any publication this work should be appropriately cited, and the papers it cites should also be recognized. This document provides a citation for the relevant parts of the database, and also provides a reference list for the research cited by those parts.

Please note that the database version for the citations given in GtoPdb are to the most recent preceding version in which the family or its subfamilies and targets were substantially changed. The links below are to the current version. If you need to consult the cited version, rather than the most recent version, please contact the GtoPdb curators.

Database links

[SLC22 family of organic cation and anion transporters](#)

<http://www.guidetopharmacology.org/GRAC/FamilyDisplayForward?familyId=146>

[Organic cation transporters \(OCT\)](#)

<http://www.guidetopharmacology.org/GRAC/FamilyDisplayForward?familyId=196>

Transporters

[OCT1\(Organic cation transporter 1\)](#)

<http://www.guidetopharmacology.org/GRAC/ObjectDisplayForward?objectId=1019>

[OCT2\(Organic cation transporter 2\)](#)

<http://www.guidetopharmacology.org/GRAC/ObjectDisplayForward?objectId=1020>

OCT3(Organic cation transporter 3)

<http://www.guidetopharmacology.org/GRAC/ObjectDisplayForward?objectId=1021>

Organic zwitterions/cation transporters (OCTN)

<http://www.guidetopharmacology.org/GRAC/FamilyDisplayForward?familyId=197>

Transporters

OCTN1(Organic cation/carnitine transporter 1)

<http://www.guidetopharmacology.org/GRAC/ObjectDisplayForward?objectId=1022>

OCTN2(Organic cation/carnitine transporter 2)

<http://www.guidetopharmacology.org/GRAC/ObjectDisplayForward?objectId=1023>

CT2(Carnitine transporter 2)

<http://www.guidetopharmacology.org/GRAC/ObjectDisplayForward?objectId=1024>

Organic anion transporters (OATs)

<http://www.guidetopharmacology.org/GRAC/FamilyDisplayForward?familyId=198>

Transporters

OAT1(Organic anion transporter 1)

<http://www.guidetopharmacology.org/GRAC/ObjectDisplayForward?objectId=1025>

OAT2(Organic anion transporter 2)

<http://www.guidetopharmacology.org/GRAC/ObjectDisplayForward?objectId=1026>

OAT3(Organic anion transporter 3)

<http://www.guidetopharmacology.org/GRAC/ObjectDisplayForward?objectId=1027>

Organic anion transporter 4

<http://www.guidetopharmacology.org/GRAC/ObjectDisplayForward?objectId=1030>

OAT5(Organic anion transporter 5)

<http://www.guidetopharmacology.org/GRAC/ObjectDisplayForward?objectId=1029>

OAT4(Organic anion transporter 7)

<http://www.guidetopharmacology.org/GRAC/ObjectDisplayForward?objectId=1028>

Urate transporter

<http://www.guidetopharmacology.org/GRAC/FamilyDisplayForward?familyId=199>

Transporters

URAT1(Urate anion exchanger 1)

<http://www.guidetopharmacology.org/GRAC/ObjectDisplayForward?objectId=1031>

Orphan or poorly characterized SLC22 family members

<http://www.guidetopharmacology.org/GRAC/FamilyDisplayForward?familyId=200>

Transporters

ORCTL3(Organic cation transporter-like 3)

<http://www.guidetopharmacology.org/GRAC/ObjectDisplayForward?objectId=1032>

ORCTL4(Organic cation transporter-like 4)

<http://www.guidetopharmacology.org/GRAC/ObjectDisplayForward?objectId=1033>

FLIPT1(Fly-like putative transporter 1)

<http://www.guidetopharmacology.org/GRAC/ObjectDisplayForward?objectId=1034>

BOIT(Brain-type organic cation transporter)

<http://www.guidetopharmacology.org/GRAC/ObjectDisplayForward?objectId=1035>

ORCTL2(Organic cation transporter-like 2)

<http://www.guidetopharmacology.org/GRAC/ObjectDisplayForward?objectId=1036>

OAT6

<http://www.guidetopharmacology.org/GRAC/ObjectDisplayForward?objectId=1037>

SLC22A23

<http://www.guidetopharmacology.org/GRAC/ObjectDisplayForward?objectId=1038>

SLC22A24

<http://www.guidetopharmacology.org/GRAC/ObjectDisplayForward?objectId=1039>

UST6

<http://www.guidetopharmacology.org/GRAC/ObjectDisplayForward?objectId=1040>

solute carrier family 22 member 31

<http://www.guidetopharmacology.org/GRAC/ObjectDisplayForward?objectId=2557>

Atypical SLC22B subfamily

<http://www.guidetopharmacology.org/GRAC/FamilyDisplayForward?familyId=859>

Transporters

synaptic vesicle glycoprotein 2A

<http://www.guidetopharmacology.org/GRAC/ObjectDisplayForward?objectId=2634>

References

1. Burns CM and Wortmann RL. (2011) Gout therapeutics: new drugs for an old disease. *Lancet* **377**: 165-77 [PMID:20719377]
2. Cha SH, Sekine T, Kusuhara H, Yu E, Kim JY, Kim DK, Sugiyama Y, Kanai Y and Endou H. (2000) Molecular cloning and characterization of multispecific organic anion transporter 4 expressed in the placenta. *J. Biol. Chem.* **275**: 4507-12 [PMID:10660625]
3. Diaz-Torné C, Perez-Herrero N and Perez-Ruiz F. (2015) New medications in development for the treatment of hyperuricemia of gout. *Curr Opin Rheumatol* **27**: 164-9 [PMID:25603039]
4. Enomoto A, Kimura H, Chairoungdua A, Shigeta Y, Jutabha P, Cha SH, Hosoyamada M, Takeda M, Sekine T and Igarashi T *et al.*. (2002) Molecular identification of a renal urate anion exchanger that regulates blood urate levels. *Nature* **417**: 447-52 [PMID:12024214]
5. Gorboulev V, Ulzheimer JC, Akhoundova A, Ulzheimer-Teuber I, Karbach U, Quester S, Baumann C, Lang F, Busch AE and Koepsell H. (1997) Cloning and characterization of two human polyspecific organic cation transporters. *DNA Cell Biol.* **16**: 871-81 [PMID:9260930]
6. Gründemann D, Liebich G, Kiefer N, Köster S and Schömig E. (1999) Selective substrates for non-neuronal monoamine transporters. *Mol. Pharmacol.* **56**: 1-10 [PMID:10385678]
7. Gründemann D, Schechinger B, Rappold GA and Schömig E. (1998) Molecular identification of the corticosterone-sensitive extraneuronal catecholamine transporter. *Nat. Neurosci.* **1**: 349-51 [PMID:10196521]
8. Hamada T, Ichida K, Hosoyamada M, Mizuta E, Yanagihara K, Sonoyama K, Sugihara S, Igawa O, Hosoya T and Ohtahara A *et al.*. (2008) Uricosuric action of losartan via the inhibition of urate transporter 1 (URAT 1) in hypertensive patients. *Am. J. Hypertens.* **21**: 1157-62 [PMID:18670416]
9. Ichida K, Hosoyamada M, Kimura H, Takeda M, Utsunomiya Y, Hosoya T and Endou H. (2003) Urate transport via human PAH transporter hOAT1 and its gene structure. *Kidney Int.* **63**: 143-55 [PMID:12472777]
10. Kenda BM, Matagne AC, Talaga PE, Pasau PM, Differding E, Lallemand BI, Frycia AM, Moureau FG, Klitgaard HV and Gillard MR *et al.*. (2004) Discovery of 4-substituted pyrrolidone butanamides as new agents with significant antiepileptic activity. *J. Med. Chem.* **47**: 530-49 [PMID:14736235]
11. Kimura H, Takeda M, Narikawa S, Enomoto A, Ichida K and Endou H. (2002) Human organic anion transporters and human organic cation transporters mediate renal transport of prostaglandins. *J. Pharmacol. Exp. Ther.* **301**: 293-8 [PMID:11907186]
12. Klitgaard H and Verdu P. (2007) Levetiracetam: the first SV2A ligand for the treatment of epilepsy. *Expert Opin Drug Discov* **2**: 1537-45 [PMID:23484603]
13. Kusuhara H, Sekine T, Utsunomiya-Tate N, Tsuda M, Kojima R, Cha SH, Sugiyama Y, Kanai Y and Endou H. (1999) Molecular cloning and characterization of a new multispecific organic anion transporter from rat brain. *J. Biol. Chem.* **274**: 13675-80 [PMID:10224140]
14. Longo N, Frigeni M and Pasquali M. (2016) Carnitine transport and fatty acid oxidation. *Biochim. Biophys. Acta* **1863**: 2422-35 [PMID:26828774]
15. Löscher W, Gillard M, Sands ZA, Kaminski RM and Klitgaard H. (2016) Synaptic Vesicle Glycoprotein 2A

- Ligands in the Treatment of Epilepsy and Beyond. *CNS Drugs* **30**: 1055-1077 [PMID:27752944]
16. Mendoza-Torreblanca JG, Vanoye-Carlo A, Phillips-Farfán BV, Carmona-Aparicio L and Gómez-Lira G. (2013) Synaptic vesicle protein 2A: basic facts and role in synaptic function. *Eur. J. Neurosci.* **38**: 3529-39 [PMID:24102679]
 17. Nicolas JM, Hannestad J, Holden D, Kervyn S, Nabulsi N, Tytgat D, Huang Y, Chanteux H, Staelens L and Matagne A *et al.*. (2016) Brivaracetam, a selective high-affinity synaptic vesicle protein 2A (SV2A) ligand with preclinical evidence of high brain permeability and fast onset of action. *Epilepsia* **57**: 201-9 [PMID:26663401]
 18. Noyer M, Gillard M, Matagne A, Hénichart JP and Wülfert E. (1995) The novel antiepileptic drug levetiracetam (ucb L059) appears to act via a specific binding site in CNS membranes. *Eur. J. Pharmacol.* **286**: 137-46 [PMID:8605950]
 19. Pao SS, Paulsen IT and Saier Jr MH. (1998) Major facilitator superfamily *Microbiol. Mol. Biol. Rev.* **62**: 1-34 [PMID:9529885]
 20. Perland E, Bagchi S, Klaesson A and Fredriksson R. (2017) Characteristics of 29 novel atypical solute carriers of major facilitator superfamily type: evolutionary conservation, predicted structure and neuronal co-expression. *Open Biol* **7**: [PMID:28878041]
 21. Perland E and Fredriksson R. (2017) Classification Systems of Secondary Active Transporters. *Trends Pharmacol. Sci.* **38**: 305-315 [PMID:27939446]
 22. Reddy VS, Shlykov MA, Castillo R, Sun EI and Saier Jr MH. (2012) The major facilitator superfamily (MFS) revisited. *FEBS J.* **279**: 2022-35 [PMID:22458847]
 23. Shu Y, Sheardown SA, Brown C, Owen RP, Zhang S, Castro RA, Ianculescu AG, Yue L, Lo JC and Burchard EG *et al.*. (2007) Effect of genetic variation in the organic cation transporter 1 (OCT1) on metformin action. *J. Clin. Invest.* **117**: 1422-31 [PMID:17476361]
 24. Takeda M, Babu E, Narikawa S and Endou H. (2002) Interaction of human organic anion transporters with various cephalosporin antibiotics. *Eur. J. Pharmacol.* **438**: 137-42 [PMID:11909604]
 25. Vogl C, Tanifuji S, Danis B, Daniels V, Foerch P, Wolff C, Whalley BJ, Mochida S and Stephens GJ. (2015) Synaptic vesicle glycoprotein 2A modulates vesicular release and calcium channel function at peripheral sympathetic synapses. *Eur. J. Neurosci.* **41**: 398-409 [PMID:25484265]
 26. von Rosenstiel P. (2007) Brivaracetam (UCB 34714). *Neurotherapeutics* **4**: 84-7 [PMID:17199019]
 27. Youngblood GL and Sweet DH. (2004) Identification and functional assessment of the novel murine organic anion transporter Oat5 (Slc22a19) expressed in kidney. *Am. J. Physiol. Renal Physiol.* **287**: F236-44 [PMID:15068970]
 28. Yu Z, Fong WP and Cheng CH. (2007) Morin (3,5,7,2',4'-pentahydroxyflavone) exhibits potent inhibitory actions on urate transport by the human urate anion transporter (hURAT1) expressed in human embryonic kidney cells. *Drug Metab. Dispos.* **35**: 981-6 [PMID:17325024]
 29. Zhang L, Schaner ME and Giacomini KM. (1998) Functional characterization of an organic cation transporter (hOCT1) in a transiently transfected human cell line (HeLa). *J. Pharmacol. Exp. Ther.* **286**: 354-61 [PMID:9655880]
 30. Zhu HJ, Appel DI, Gründemann D and Markowitz JS. (2010) Interaction of organic cation transporter 3 (SLC22A3) and amphetamine. *J. Neurochem.* **114**: 142-9 [PMID:20402963]