

SLC47 family of multidrug and toxin extrusion transporters in GtoPdb v.2021.3

Ken-ichi Inui¹

1. Kyoto Pharmaceutical University, Japan

Abstract

These proton:organic cation exchangers are predicted to have 13 TM segments [12] and are suggested to be responsible for excretion of many drugs in the liver and kidneys [1].

Contents

This is a citation summary for SLC47 family of multidrug and toxin extrusion transporters in the [Guide to Pharmacology](#) database (GtoPdb). It exists purely as an adjunct to the database to facilitate the recognition of citations to and from the database by citation analyzers. Readers will almost certainly want to visit the relevant sections of the database which are given here under database links.

GtoPdb is an expert-driven guide to pharmacological targets and the substances that act on them. GtoPdb is a reference work which is most usefully represented as an on-line database. As in any publication this work should be appropriately cited, and the papers it cites should also be recognized. This document provides a citation for the relevant parts of the database, and also provides a reference list for the research cited by those parts. For further details see [2].

Please note that the database version for the citations given in GtoPdb are to the most recent preceding version in which the family or its subfamilies and targets were substantially changed. The links below are to the current version. If you need to consult the cited version, rather than the most recent version, please contact the GtoPdb curators.

Database links

SLC47 family of multidrug and toxin extrusion transporters

<https://www.guidetopharmacology.org/GRAC/FamilyDisplayForward?familyId=236>

Transporters

[MATE1\(Multidrug and toxin extrusion\)](https://www.guidetopharmacology.org/GRAC/ObjectDisplayForward?objectId=1216)

<https://www.guidetopharmacology.org/GRAC/ObjectDisplayForward?objectId=1216>

[MATE2-K\(MATE2\)](https://www.guidetopharmacology.org/GRAC/ObjectDisplayForward?objectId=1217)

<https://www.guidetopharmacology.org/GRAC/ObjectDisplayForward?objectId=1217>

References

1. Alexander SPH, Kelly E, Mathie A, Peters JA, Veale EL, Armstrong JF, Faccenda E, Harding SD, Pawson AJ and Sharman JL *et al.*. (2019) THE CONCISE GUIDE TO PHARMACOLOGY 2019/20: Transporters. *Br J Pharmacol* **176 Suppl 1**: S397-S493 [\[PMID:31710713\]](#)
2. Buneman P, Christie G, Davies JA, Dimitrellou R, Harding SD, Pawson AJ, Sharman JL and Wu Y. (2020) Why data citation isn't working, and what to do about it *Database* **2020** [\[PMID:32367113\]](#)
3. Chen Y, Zhang S, Sorani M and Giacomini KM. (2007) Transport of paraquat by human organic cation transporters and multidrug and toxic compound extrusion family. *J Pharmacol Exp Ther* **322**: 695-700 [\[PMID:17495125\]](#)
4. Ito S, Kusuhara H, Kuroiwa Y, Wu C, Moriyama Y, Inoue K, Kondo T, Yuasa H, Nakayama H and Horita S *et al.*. (2010) Potent and specific inhibition of mMate1-mediated efflux of type I organic cations in the liver and kidney by pyrimethamine. *J Pharmacol Exp Ther* **333**: 341-50

[PMID:20065018]

5. Masuda S, Terada T, Yonezawa A, Tanihara Y, Kishimoto K, Katsura T, Ogawa O and Inui K. (2006) Identification and functional characterization of a new human kidney-specific H+/organic cation antiporter, kidney-specific multidrug and toxin extrusion 2. *J Am Soc Nephrol* **17**: 2127-35 [PMID:16807400]
6. Ohta KY, Inoue K, Hayashi Y and Yuasa H. (2006) Molecular identification and functional characterization of rat multidrug and toxin extrusion type transporter 1 as an organic cation/H+ antiporter in the kidney. *Drug Metab Dispos* **34**: 1868-74 [PMID:16928787]
7. Otsuka M, Matsumoto T, Morimoto R, Arioka S, Omote H and Moriyama Y. (2005) A human transporter protein that mediates the final excretion step for toxic organic cations. *Proc Natl Acad Sci USA* **102**: 17923-8 [PMID:16330770]
8. Tanihara Y, Masuda S, Sato T, Katsura T, Ogawa O and Inui K. (2007) Substrate specificity of MATE1 and MATE2-K, human multidrug and toxin extrusions/H(+) - organic cation antiporters. *Biochem Pharmacol* **74**: 359-71 [PMID:17509534]
9. Terada T, Masuda S, Asaka J, Tsuda M, Katsura T and Inui K. (2006) Molecular cloning, functional characterization and tissue distribution of rat H+/organic cation antiporter MATE1. *Pharm Res* **23**: 1696-701 [PMID:16850272]
10. Tsuda M, Terada T, Ueba M, Sato T, Masuda S, Katsura T and Inui K. (2009) Involvement of human multidrug and toxin extrusion 1 in the drug interaction between cimetidine and metformin in renal epithelial cells. *J Pharmacol Exp Ther* **329**: 185-91 [PMID:19164462]
11. Yasujima T, Ohta KY, Inoue K, Ishimaru M and Yuasa H. (2010) Evaluation of 4',6-diamidino-2-phenylindole as a fluorescent probe substrate for rapid assays of the functionality of human multidrug and toxin extrusion proteins. *Drug Metab Dispos* **38**: 715-21 [PMID:20047987]
12. Zhang X and Wright SH. (2009) MATE1 has an external COOH terminus, consistent with a 13-helix topology. *Am J Physiol Renal Physiol* **297**: F263-71 [PMID:19515813]