

Succinate receptor (version 2019.4) in the IUPHAR/BPS Guide to Pharmacology Database

Anthony P. Davenport¹, Julien Hanson² and Wen Chiy Liew³

1. University of Cambridge, UK
2. University of Liege, Belgium
3. University of Edinburgh, UK

Abstract

Nomenclature as recommended by NC-IUPHAR [7]. The Succinate receptor has been identified as being activated by physiological levels of the Krebs's cycle intermediate succinate and other dicarboxylic acids such as maleate in 2004. Since its pairing with its endogenous ligand, the receptor has been the focus of intensive research and its role has been evidenced in various (patho)physiological processes such as regulation of renin production, retinal angiogenesis, inflammation or immune response.

Contents

This is a citation summary for Succinate receptor in the [Guide to Pharmacology](#) database (GtoPdb). It exists purely as an adjunct to the database to facilitate the recognition of citations to and from the database by citation analyzers. Readers will almost certainly want to visit the relevant sections of the database which are given here under database links.

[GtoPdb](#) is an expert-driven guide to pharmacological targets and the substances that act on them. GtoPdb is a reference work which is most usefully represented as an on-line database. As in any publication this work should be appropriately cited, and the papers it cites should also be recognized. This document provides a citation for the relevant parts of the database, and also provides a reference list for the research cited by those parts.

Please note that the database version for the citations given in GtoPdb are to the most recent preceding version in which the family or its subfamilies and targets were substantially changed. The links below are to the current version. If you need to consult the cited version, rather than the most recent version, please contact the GtoPdb curators.

Database links

[Succinate receptor](#)

<http://www.guidetopharmacology.org/GRAC/FamilyDisplayForward?familyId=446>

[Introduction to Succinate receptor](#)

<http://www.guidetopharmacology.org/GRAC/FamilyIntroductionForward?familyId=446>

Receptors

[succinate receptor](#)

References

1. Abbracchio MP, Burnstock G, Boeynaems JM, Barnard EA, Boyer JL, Kennedy C, Knight GE, Fumagalli M, Gachet C and Jacobson KA *et al.*. (2006) International Union of Pharmacology LVIII: update on the P2Y G protein-coupled nucleotide receptors: from molecular mechanisms and pathophysiology to therapy. *Pharmacol. Rev.* **58**: 281-341 [PMID:16968944]
2. Aguiar CJ, Andrade VL, Gomes ER, Alves MN, Ladeira MS, Pinheiro AC, Gomes DA, Almeida AP, Goes AM and Resende RR *et al.*. (2010) Succinate modulates Ca(2+) transient and cardiomyocyte viability through PKA-dependent pathway. *Cell Calcium* **47**: 37-46 [PMID:20018372]
3. Ariza AC, Deen PM and Robben JH. (2012) The succinate receptor as a novel therapeutic target for oxidative and metabolic stress-related conditions. *Front Endocrinol (Lausanne)* **3**: 22 [PMID:22649411]
4. Bhuniya D, Umrani D, Dave B, Salunke D, Kukreja G, Gundu J, Naykodi M, Shaikh NS, Shitole P and Kurhade S *et al.*. (2011) Discovery of a potent and selective small molecule hGPR91 antagonist *Bioorg. Med. Chem. Lett.* **21**: 3596-602 [PMID:21571530]
5. Cantagrel V, Lossi AM, Boulanger S, Depetris D, Mattei MG, Gecz J, Schwartz CE, Van Maldergem L and Villard L. (2004) Disruption of a new X linked gene highly expressed in brain in a family with two mentally retarded males. *J. Med. Genet.* **41**: 736-42 [PMID:15466006]
6. Correa PR, Kruglov EA, Thompson M, Leite MF, Dranoff JA and Nathanson MH. (2007) Succinate is a paracrine signal for liver damage. *J. Hepatol.* **47**: 262-9 [PMID:17451837]
7. Davenport AP, Alexander SP, Sharman JL, Pawson AJ, Benson HE, Monaghan AE, Liew WC, Mpamhanga CP, Bonner TI and Neubig RR *et al.*. (2013) International Union of Basic and Clinical Pharmacology. LXXXVIII. G protein-coupled receptor list: recommendations for new pairings with cognate ligands. *Pharmacol. Rev.* **65**: 967-86 [PMID:23686350]
8. Fredriksson R, Lagerström MC, Lundin LG and Schiöth HB. (2003) The G-protein-coupled receptors in the human genome form five main families. Phylogenetic analysis, paralogon groups, and fingerprints. *Mol. Pharmacol.* **63**: 1256-72 [PMID:12761335]
9. Geubelle P, Gilissen J, Dilly S, Poma L, Dupuis N, Laschet C, Abboud D, Inoue A, Jouret F and Pirote B *et al.*. (2017) Identification and pharmacological characterization of succinate receptor agonists. *Br. J. Pharmacol.* **174**: 796-808 [PMID:28160606]
10. Gilissen J, Geubelle P, Dupuis N, Laschet C, Pirote B and Hanson J. (2015) Forskolin-free cAMP assay for Gi-coupled receptors. *Biochem. Pharmacol.* **98**: 381-91 [PMID:26386312]
11. Gnana-Prakasam JP, Ananth S, Prasad PD, Zhang M, Atherton SS, Martin PM, Smith SB and Ganapathy V. (2011) Expression and iron-dependent regulation of succinate receptor GPR91 in retinal pigment epithelium. *Invest. Ophthalmol. Vis. Sci.* **52**: 3751-8 [PMID:21357408]
12. Gonzalez NS, Communi D, Hannedouche S and Boeynaems JM. (2004) The fate of P2Y-related orphan receptors: GPR80/99 and GPR91 are receptors of dicarboxylic acids. *Purinergic Signal.* **1**: 17-20 [PMID:18404396]
13. Hakak Y, Lehmann-Bruinsma K, Phillips S, Le T, Liaw C, Connolly DT and Behan DP. (2009) The role of the GPR91 ligand succinate in hematopoiesis. *J. Leukoc. Biol.* **85**: 837-43 [PMID:19204147]
14. He W, Miao FJ, Lin DC, Schwandner RT, Wang Z, Gao J, Chen JL, Tian H and Ling L. (2004) Citric acid cycle intermediates as ligands for orphan G-protein-coupled receptors. *Nature* **429**: 188-93 [PMID:15141213]
15. Hebert SC. (2004) Physiology: orphan detectors of metabolism. *Nature* **429**: 143-5 [PMID:15141197]
16. Högberg C, Gidlöf O, Tan C, Svensson S, Nilsson-Öhman J, Erlinge D and Olde B. (2011) Succinate independently stimulates full platelet activation via cAMP and phosphoinositide 3-kinase-β signaling. *J. Thromb. Haemost.* **9**: 361-72 [PMID:21143371]
17. Joost P and Methner A. (2002) Phylogenetic analysis of 277 human G-protein-coupled receptors as a tool for the prediction of orphan receptor ligands. *Genome Biol.* **3**: RESEARCH0063 [PMID:12429062]
18. Littlewood-Evans A, Sarret S, Apfel V, Loesle P, Dawson J, Zhang J, Muller A, Tigani B, Kneuer R and

- Patel S *et al.*. (2016) GPR91 senses extracellular succinate released from inflammatory macrophages and exacerbates rheumatoid arthritis. *J. Exp. Med.* **213**: 1655-62 [PMID:27481132]
19. Macaulay IC, Tijssen MR, Thijssen-Timmer DC, Gusnanto A, Steward M, Burns P, Langford CF, Ellis PD, Dudbridge F, Zwaginga JJ, Watkins NA, van der Schoot CE and Ouwehand WH. (2007) Comparative gene expression profiling of in vitro differentiated megakaryocytes and erythroblasts identifies novel activatory and inhibitory platelet membrane proteins. *Blood* **109**: 3260-9 [PMID:17192395]
 20. Molnár T, Héja L, Emri Z, Simon A, Nyitrai G, Pál I and Kardos J. (2011) Activation of astroglial calcium signaling by endogenous metabolites succinate and gamma-hydroxybutyrate in the nucleus accumbens. *Front Neuroenergetics* **3**: 7 [PMID:22180742]
 21. Regard JB, Sato IT and Coughlin SR. (2008) Anatomical profiling of G protein-coupled receptor expression. *Cell* **135**: 561-71 [PMID:18984166]
 22. Rubic T, Lametschwandner G, Jost S, Hinteregger S, Kund J, Carballido-Perrig N, Schwärzler C, Junt T, Voshol H and Meingassner JG *et al.*. (2008) Triggering the succinate receptor GPR91 on dendritic cells enhances immunity. *Nat. Immunol.* **9**: 1261-9 [PMID:18820681]
 23. Rubić-Schneider T, Carballido-Perrig N, Regairaz C, Raad L, Jost S, Rauld C, Christen B, Wieczorek G, Kreutzer R and Dawson J *et al.*. (2017) GPR91 deficiency exacerbates allergic contact dermatitis while reducing arthritic disease in mice. *Allergy* **72**: 444-452 [PMID:27527650]
 24. Sapieha P, Sirinyan M, Hamel D, Zaniolo K, Joyal JS, Cho JH, Honoré JC, Kermorvant-Duchemin E, Varma DR and Tremblay S *et al.*. (2008) The succinate receptor GPR91 in neurons has a major role in retinal angiogenesis. *Nat. Med.* **14**: 1067-76 [PMID:18836459]
 25. Southern C, Cook JM, Neetoo-Isseljee Z, Taylor DL, Kettleborough CA, Merritt A, Bassoni DL, Raab WJ, Quinn E and Wehrman TS *et al.*. (2013) Screening β -Arrestin Recruitment for the Identification of Natural Ligands for Orphan G-Protein-Coupled Receptors. *J Biomol Screen* **18**: 599-609 [PMID:23396314]
 26. Sundström L, Greasley PJ, Engberg S, Wallander M and Ryberg E. (2013) Succinate receptor GPR91, a $G\alpha(i)$ coupled receptor that increases intracellular calcium concentrations through PLC β . *FEBS Lett.* **587**: 2399-404 [PMID:23770096]
 27. Toma I, Kang JJ, Sipos A, Vargas S, Bansal E, Hanner F, Meer E and Peti-Peterdi J. (2008) Succinate receptor GPR91 provides a direct link between high glucose levels and renin release in murine and rabbit kidney. *J. Clin. Invest.* **118**: 2526-34 [PMID:18535668]
 28. Vargas SL, Toma I, Kang JJ, Meer EJ and Peti-Peterdi J. (2009) Activation of the succinate receptor GPR91 in macula densa cells causes renin release. *J. Am. Soc. Nephrol.* **20**: 1002-11 [PMID:19389848]
 29. Wittenberger T, Schaller HC and Hellebrand S. (2001) An expressed sequence tag (EST) data mining strategy succeeding in the discovery of new G-protein coupled receptors. *J. Mol. Biol.* **307**: 799-813 [PMID:11273702]
 30. Zhang D, Gao ZG, Zhang K, Kiselev E, Crane S, Wang J, Paoletta S, Yi C, Ma L and Zhang W *et al.*. (2015) Two disparate ligand-binding sites in the human P2Y1 receptor. *Nature* **520**: 317-21 [PMID:25822790]
 31. Zhang J, Zhang K, Gao ZG, Paoletta S, Zhang D, Han GW, Li T, Ma L, Zhang W and Müller CE *et al.*. (2014) Agonist-bound structure of the human P2Y12 receptor. *Nature* **509**: 119-22 [PMID:24784220]
 32. Zhang K, Zhang J, Gao ZG, Zhang D, Zhu L, Han GW, Moss SM, Paoletta S, Kiselev E and Lu W *et al.*. (2014) Structure of the human P2Y12 receptor in complex with an antithrombotic drug. *Nature* **509**: 115-8 [PMID:24670650]