

---

# Bipolar flaking as a component of a supraregional lithic resource base: A comparative study of cores from the Pampean and Northcentral Patagonian Atlantic coasts (Argentina)

Anahí Banegas<sup>1</sup>, Mariano Bonomo<sup>2</sup>, Julieta Gómez Otero<sup>1</sup>

1. CONICET - Instituto de Diversidad y Evolución Austral. B. Brown 2915, Puerto Madryn, Argentina.

Email: Banegas: banegas.anahi@yahoo.com.ar; Gómez Otero: julietagomezotero@yahoo.com.ar

2. CONICET - División Arqueología, Facultad de Ciencias Naturales y Museo, Universidad Nacional de La Plata, La Plata, Argentina. Email: mbonomo@fcnym.unlp.edu.ar

---

## Abstract:

Several authors argue that bipolar technology is an expeditious and versatile strategy, efficient for the reduction of small nodules. However, few studies analyze the use of bipolar reduction in relation to the lithic environmental supply. At the Pampean and Patagonian maritime coast, the technique has been widely recorded and seems to be conspicuous in indigenous populations that occupied coastal areas. With the aim of assessing and comparing the application of bipolar reduction 196 nuclei recovered from sites of diverse chronologies in the Pampean and northcentral Patagonian coasts were analyzed. The use of the bipolar technique was observed in relation to the lithic environmental structure, the size of nodules and their raw material. The results revealed high frequencies of bipolar nuclei at the Pampean coast (87%) and moderate frequencies at the northcentral Patagonian coast (39%). It was determined that the variables that influenced the application of bipolar techniques the most were the small size of the nodules and their rounded shape, generally thick, and without flat surfaces. As for raw material quality, the technique was applied on good to medium quality rocks at the Pampean coast, while in the northcentral coast it was exclusively applied on pebbles and cobbles of superior qualities (very good to excellent). The differences observed are most likely linked with technological choices made by different coastal populations to make the best use of widely available local rocks.

**Keywords:** bipolar flaking; lithic resource base; cores; comparative study; Pampean and northcentral Patagonian Atlantic coasts

## 1. Introduction

Bipolar flaking on an anvil is one of the earliest lithic reduction techniques in the history of humankind. It has been documented in diverse archaeological contexts from all over the world and covers a wide timespan (Arthur 2010; Le Brun-Ricalens 2005; Guyodo & Marchand 2005; Nami 2000; Pargeter & Metin 2017; Shott 1989; Soriano *et al.* 2011). It has



also been documented by ethnographers and ethnoarchaeologists among diverse populations using traditional technologies (Flood 1980; Hayden 1980; Sillitoe & Hardy 2003; White 1968). Intense debates have taken place regarding the aims for which human groups applied the technique, attributed both to the physical and natural constraints of lithic resources and to social factors related to stylistic preferences and cultural traditions or ethnicities (Flegenheimer *et al.* 1995). It has recently been defined as a sophisticated technique that demands continuous decision-making, intensive learning and practice, and may be associated with women (Arthur 2010). Frequently mentioned among the main functional causes behind its application are the use of rock supports too small or of poor flaking quality (Binford & Quimby 1963; Flenniken 1981); the preservation or recycling of raw material of high quality or from distant sources (Goodyear 1993) and the optimization of the time and energy invested in tool manufacture (Nami 2000; Shott 1989). Its use has also been suggested in the manufacturing of microliths to insert in compound tools such as Australian spears (Flegenheimer *et al.* 1995; Flood 1980; Goodyear 1993; Hayden 1980). Whatever these causes may be, there is common agreement among archaeologists across the globe that the bipolar technique is versatile and within the range of expedient strategies that favors an efficient use of nodules.

The naturalist Florentino Ameghino (1910) recorded it in the Atlantic coast of Argentina at the beginning of the 20<sup>th</sup> century, and deemed it a simple and “primitive” procedure applied by a predecessor of modern humans, the *Homo pampæus*, to manufacture stone tools by breaking rounded beach-deposited pebbles. Some authors assigned it to coastal populations with economies oriented towards the exploitation of marine resources in the Pampean and Patagonian regions (Ameghino 1910; Bórmida 1964; Conlazo 1983; Díaz de Chiri 1977; Menghin 1952 1963) (Figure 1). Others suggested links with hunter-gatherer groups from the inland plains that, either occasionally or sporadically, occupied the littoral zone and made use of the available lithic and faunal resources (Aparicio 1932; Holmes 1912; Hrdlička 1912; Outes 1909; Politis 1984). Based on an Austro-German culture-historical perspective, Menghin and Bórmida (Bórmida 1964; Menghin 1952) suggested that bipolar flaking corresponded to an ancient “epiprotolithic” tradition of inferior coastal foragers that made use of the rounded pebbles available at the Pampean and Patagonian coastlines to manufacture choppers, unifacial tools, scrapers and denticulates. They distinguished this tradition from a more modern one, named “miolithic”, of superior inland hunters who produced flakes, blades and bifacial tools like projectile points and made eventual use of coastal areas. Towards the late 1980s, new theoretical and methodological approaches derived from processual approach enabled archaeologists to refute and overcome Menghin and Bórmida's main assertions (Borrero 2015; Cruz & Caracotche 2008; Gómez Otero 1996; Politis 1984; Orquera & Gómez Otero 2008). In this regard, Bonomo (2005) was able to demonstrate that the lithic workshops with abundant bipolar artefacts from the dunes along the coast of the *Interserrana* area of the Pampean region were produced by hunter-gatherers occupying both the Atlantic coast and the inland plains (Figure 1). Subsequent advances in Pampean and Patagonian archaeology pointed to a significant spatial and temporal variability in the use of coastal areas and a diversity of technological strategies used by past human populations (Orquera & Gómez Otero 2008).

A comparative technological study of lithic assemblages from the coast of the *Interserrana* area of the Pampean region and the northcentral Patagonian coast carried out by Banegas (2016) enabled the observation of differences and similarities between both sectors (Figure 1). One of the technological features shared by both areas is the bipolar flaking of pebbles and cobbles (Banegas 2016: 240). This work presents the results of the comparative analysis of 196 cores - bipolar and not - recovered from 14 archaeological sites of diverse chronology along the Pampean coast (between Cape Corrientes and the Quequén Salado river

mouth) and the northcentral Patagonian coast (between Verde Stream and Cracker Bay) (Figure 1). The aim is to assess the application of bipolar flaking in both areas. Spatial and temporal trends are explored and compared with the archaeological evidence of the bipolar technique in other sectors of the continental Atlantic coast. We have chosen to analyze only the cores from among all of the recovered artefactual classes, as these provide the most information about the first stages of lithic raw material exploitation for tool manufacturing, that is to say, the initial selection and initial reduction of the nodules. Diagnostic features of bipolar technique can be found in both cores and flakes: for example, chipped or splintered butts at opposing ends, with columnar extractions, flat flakes without or with negative bulbs, among others (Binford & Quimby 1963; Curtoni 1996; Kobayashi 1975; Nami 2000).




Figure 1. A: Study area and A.1 and A.2 archaeological sites mentioned in the text.

### 1.1. Areas of study and research background

The Pampean coastal sector under study lies within the *Interserrana* area, Buenos Aires province, and extends from Cape Corrientes to the mouth of the Quequén Salado river (Figure 1). Two geomorphological sectors can be distinguished: a higher one with cliffs carved by marine erosion, and a lower one characterized by a dune barrier formed by the deposition of sandy sediments (Bonomo 2005: 36-39). The coastal gravel is mainly comprised of clasts of volcanic origin carried by coastal drift currents and deposited on the beaches of Buenos Aires province. Nowadays these deposits are mainly located within the area under study, but very scarce quantities can also be found north of Cape Corrientes and southwest of the Quequén Salado river (Bayón & Zavala 1997).

Archaeological research shows that since the early to mid-Holocene, human populations intensively exploited this coastal environment and its marine resources (Bayón *et al.* 2012; Bonomo & León 2010; Bonomo *et al.* 2013; Politis & Bayón 1996; Politis *et al.* 2009). However, earlier evidence of coastal pebbles and cobbles, and marine molluscs has been found at archaeological sites of the inland plains, indicating the Pampean coast was exploited since the beginning of the human occupation of the region, that took place during the late Pleistocene (Bonomo 2005: 263). In the coastal lithic assemblages, debitage and cores predominate, while tools manufactured by flaking (such as end-scrapers, side-scrapers, and knives), anvils and hammerstones appear in lesser quantities. The most commonly used raw materials are basalt pebbles, although quartzites from the Sierras Bayas formation, located inland between 80 and 160 km away from the coast, and a small amount of obsidian of unknown origin were also used (Bonomo 2005). Quartzites are the most used raw material along the plains of the Pampean region since the late Pleistocene until late Holocene.

The northcentral coast of Patagonia extends from the mouth of the Verde stream to Cracker Bay (Figure 1). It is characterized by plateaus that do not exceed 100 m a.s.l., dissected by canyons and gullies (Súnico 1996). A succession of coastal bars, palaeocliffs, and ancient tidal flats are the result of isostatic and eustatic changes, and neotectonics during the Pleistocene and Holocene (Codignotto *et al.* 1992; González & Weiler 1994). Dunes have developed atop the cliffs and on the coastal bars of ancient bays (Súnico 1996). Three types of coastline can be distinguished: with gulfs (San Matías, San José and Nuevo), with spits and barrier islands at Valdés Cove and with an open sea coastline at the Valdés Peninsula. As regards geology, Tertiary and Quaternary-aged sedimentary rocks prevail (Haller 1981). The supply of lithic raw material for flaking is abundant: secondary deposits of pebbles are available, as well as blocks of ignimbrites and chalcedonies from veins of the Marifil formation (Malvicini & Llambías 1974) that outcrops at the Verde Stream.

Archaeological research shows that the area has been occupied at least since the Middle Holocene - between 7400 and 200 years BP - but with spatial and temporal variations in the use of coastal space (Gómez Otero 2006; Gómez Otero *et al.* 1999; 2013). The highest archaeological density has been determined close to the present coastline, which indicates that mobility and settlement were mostly restricted to the littoral sector (Gómez Otero 2006). A progressive increase in the amount of sites and diversity of activities at these sites has been recorded starting at 2000 BP. The diet combined terrestrial resources (*guanacos* and plants) complemented with marine resources such as molluscs, pinnipeds and fish (Gómez Otero 2006 2007; Svoboda 2015). As regards lithic technology, local pebbles of basalt and flint of very good flaking quality and, less frequently, rhyolites of medium quality, were used. Non-local rocks are represented by six obsidian varieties, among which stand out the T/SCI and SI types from sources located on the northcentral plateau 180 to 300 km away from the coast stand out (Gómez Otero & Stern 2005). Other frequently chosen rocks are chalcedonies and fossilized wood, but their origin has not yet been established (Banegas *et al.* 2016).

## 2. Materials and methods

The characterization of the regional lithic resource base was carried out by Bonomo (2005) for the Pampean coast, and Banegas, Goye and Gómez Otero (2016) for the northcentral Patagonian coast. It is worth mentioning that for northcentral Patagonian coast, the geographic space was split into two large analytical units following Gomez Otero's (2006) model: the “Valdés Peninsula” and “Outside the Valdés Peninsula” that in turn comprise diverse coastal types. Three temporal blocks were also distinguished: Middle Holocene (from 7400 to 4000 years BP); initial late Holocene (from 3900 to 1000 years BP) and final late Holocene (from 1000 to 300 years BP) (Gómez Otero 2006) (Table 1).

Table 1. Archaeological sites with their contextual information and the analysed samples. H=Holocene.

Sector	Site (total number of lithic artefacts)	Site function	Period	Total number of cores	Number of bipolar cores
Pampean coast	Bellamar 1 (1133)	lithic workshop	Middle to late H.	55	47
	Bellamar 2 (74)	lithic workshop	Middle to late H.	16	15
	La Eufemia (175)	lithic workshop	Middle to late H.	13	11
<b>Total</b>				<b>84</b>	<b>73 (87%)</b>
Valdés Peninsula (VP)	Las Lisas 2 C2-M2 (363)	location	Initial late H.	18	9
	Flechero del 39 (229)	location	Initial late H.	6	1
	Las Ollas 1 M1 (81)	residential base	Final late H.	6	4
	Pta. Cormoranes 3-F1 (11)	station	Middle H.	1	1
	San Pablo 4-M3 (101)	location	Initial late H.	2	1
	San Pablo 6-C2-M2 (61)	midden	Final late H.	1	1
<b>Total</b>				<b>34</b>	<b>17 (50%)</b>
Outside VP	Aº Verde M1-2 (214)	location	Middle H.	15	--
	R. de Elizalde 1 (411)	temporary camp	Initial late H.	28	24
	R. de Elizalde 5 (93)	location	Final late H.	4	1
	B. Cracker 4M1 (166)	location	Middle H.	27	1
	B. Cracker 8 (83)	location	Middle H.	4	1
<b>Total</b>				<b>78</b>	<b>27 (34%)</b>

The methodology applied by Bonomo (2005) on the Pampean coast consisted of a simple random sampling of the natural deposits of coastal pebbles and cobbles following 1.5 m-wide transects and 0.5 m<sup>2</sup> grids, in order to analyse their spatial distribution, topographic location in the beach's profile, maximum dimensions of clasts and most abundant raw materials. Ten years later, additional sampling was carried out on different marine geoforms from the northcentral Patagonian coast, where every pebble that could be collected within a predefined time-span of between 15 and 20 minutes was considered (Banegas *et al.* 2016: 38). The maximum dimensions, shapes and lithological classes were determined, following some of the analytical procedures applied by Berón *et al.*, (1995), Franco & Borrero (1999), Nami (1992) and Shelley (1993). To document the mineral compositions of the rocks in both areas, thirty petrographic thin sections were made from coastal gravels and lithic artefacts (8 from Pampean archaeological sites and 20 from the northcentral Patagonia coast) (Banegas 2016: 274-283; Bonomo & Prates 2014).

The 196 cores under study were collected from systematic samplings in archaeological sites of different function and chronology (Table 1). To identify bipolar cores, the presence of

the following attributes was considered: 1) unprepared, splintered or lack of striking platforms, 2) crush marks, and or hinges in one or two poles or opposing distal ends, 3) negatives of flake removals originating from a single or from opposed poles, 4) flake scars on one or both faces, that can cover the full core's original length, and 5) frequent stepped fractures. For each core, the raw material and morphological type were recorded, following different authors: Aschero (1975; 1983: 12) and Prous (2004: 54-57).

For bipolar cores the approach of Bonomo (2005: 97) was followed, which distinguishes four groups according to the degree of modification suffered by the extremes of the original pebble or cobble: a) a single modified pole with flake scars, b) both poles modified, c) with three or more poles, d) undifferentiated by fracture (Bonomo 2002). The quality of raw materials was assessed following Franco and Aragón's (2004) criteria. The metric variables - maximum length, width and thickness in mm - were measured with a digital slide calliper. To estimate the degree of core modification, they were grouped according to the proportion of surface covered by cortex, at 20% intervals, as well as distinguishing between active and depleted cores. It should be noted that the coastal pebbles' cortex has not been produced by physico-chemical changes but by marine transport. This mechanical alteration transforms the external surface of the clasts, resulting in a finer grained matrix on the outside as compared to the inside of the nodule. In other words, the attrition taking place during marine transport homogenizes the exterior surface of rocks by reducing their granulometry (Bonomo 2005: 92). This particular type of cortex, which characterizes rocks from secondary deposits, is named *néo-cortex* in the French literature (Inizan *et al.* 1995).

### 3. Data results

#### 3.1. Characterization of the regional base of lithic resources base

Pampean coast: the rock deposits comprise pebbles and cobbles measuring from 3 to 15 cm in maximum diameter. Clasts with a diameter greater than 3 cm were selected. The average maximum length of the collected rocks is of 9 cm (Bonomo & Prates 2014). As for morphology, the large majority of clasts present elliptical and flattened shapes produced by marine transport (Figure 2). Among the 191 rocks collected, andesites-dacites abound (41%), followed by rhyolites (27%), basalts (16%) and flint (8%). The rocks of the best flaking quality are the siliceous ones, while the rest of the lithologies (basalts, andesites-dacites, quartzites, chert) exhibit predominantly good to medium qualities.

Patagonian southcentral coast: the deposits comprise pebbles and cobbles with a maximum diameter of between 2.7 and 13 cm, and an average of 7 cm. It is worth noting that those sampled in VP are smaller (between 4 and 6 cm) than in the other sectors (Figure 2). Flat, rounded rocks predominate, except in Valdés Cove, where they are rounded but thick. The nodules studied from western San Matías gulf, however, present sub-rounded and angular shapes (Banegas 2016: 65-67). The most abundant raw materials are basalt (42%), rhyolite (18%) and chert (14%). The rest (andecite-dacites; ignimbrites; quartzites, chalcedony and fossilized woods) do not exceed 6.5%. Some differences were observed between these coastal environments. At the San Matías gulf area, the supply of rhyolites of medium quality stands out (34%) and there are also veins of very good to excellent quality chalcedonies (13%). On the contrary, basalts of different qualities prevail at the Valdés Peninsula (VP) and the south of the Nuevo gulf (56% and 48% respectively). Those of higher quality (good to excellent) were only documented in VP.





Figure 2. Samplings of lithic raw materials on the beaches. A) Pampean coast, B) northcentral Patagonian coast

### 3.2. Core morphological types and raw materials

Among the 84 cores from the Pampean coast under study, four morphological types were recognized: bipolar, tabular, discoidal and one with isolated flaking (Figure 3). Almost all of the raw materials modules are pebbles and cobbles (97.6%), except for two rhyolite tabular chunks. The predominant core type is bipolar ( $n = 73$ : 87%), among which prevail those cores with two modified poles ( $n = 34$ : 43.5%), followed by those with only one modified pole ( $n = 18$ : 23%) (Figure 3 and Table 2).

On the northcentral Patagonian coast, seven morphological types were identified among the 112 recovered cores. Three stand out: bipolar ( $n = 44$ : 39%), with isolated flaking ( $n = 31$ : 28%), and discoidal ( $n = 11$ : 10%) (Figure 3 and Table 2). The rest (globular, bifacial, pyramidal and polyhedral) are represented by one to five specimens and do not exceed 5% of the total. Thirteen cores were classified as undetermined ( $n = 13$ : 12%). The raw materials modules chosen are pebbles and cobbles, but also tabular chunks and undifferentiated supports. It is worth mentioning that bipolar cores reach 50% in Valdés Peninsula, and 34% outside the Valdés Peninsula.

For the sample of Pampean coast, five major lithologies were identified (andesites-dacites; basalts, quartzites, rhyolites, flint), near all of them local, except inland quartzite. Among the bipolar cores, basalts of high quality (90%) dominate, followed by very good-quality flint (4%) and medium quality andesites and medium quality rhyolites (3% of each).




Figure 3. Bipolar cores: A- core with three poles modified; B-both poles modified; C: a single modified pole.

For the sample of northcentral Patagonian coast cores, 10 lithological varieties were recognized (basalts, flint, chalcedonies, fossilized woods, obsidians, rhyolites, ignimbrites, andesites-dacites, quartzite and silicified limestones). Among those found in VP, bipolar flint cores ( $n=10$ : 59%), prevail followed by basalt and chalcedony in equal proportions ( $n= 3$ : 18% of each). The only obsidian core, of non-local origin, is bipolar. Bipolarity was not documented in fossilized wood or regular-quality rocks (“Other” categories) which comprise rhyolites, ignimbrites, andesites-dacites, quartzites and silicified limestones. Outside VP, predominate siliceous bipolar cores ( $n=17$ : 63%), followed by basalt ( $n=7$ :26%). Unlike at



VP, bipolar reduction was documented in two small fossilized wood cores (7%), which come from coarse pebbles. A single core of chalcedony (4%) was found in the Rincón de Elizalde 1 archaeological site, 1 to 20 km away from the primary source of this rock located in Verde Stream. However, the original support could not be determined because of the extensive reduction suffered by the core.

Table 2. Core morphological types.

Core morphological types	Pampean coast	Valdés Peninsula (VP)	Outside VP
Bipolar	73: 87%	17: 50%	27: 34%
Isolated flaking	1: 1%	6: 17%	25: 32%
Tabular	5: 6%	--	--
Discoidal	2: 2%	5: 15%	6: 8%
Undetermined	3: 3%	5: 15%	8: 10%
Pyramid-shaped	--	1: 3%	2: 3%
Globular	--	--	5: 6%
Bifacial	--	--	4: 5%
Polyhedral	--	--	1: 2%

### 3.3. Core size and state

On the Pampean coast small cores (74%) prevail among which 84% (n=52) are bipolar cores. In decreasing order follow medium-size cores (23%), among which bipolar cores represent a 63%. Two large cores were recorded, neither are bipolar. The minimum number of flake scars varies from one to five removals, with an average 2.98 flake scars per core. Regarding the state at the moment of discard and based on the percentage of cortex, the size and the absence of striking platforms, 54% were deemed to be depleted cores, while 32% had medium intensity modification, and 14% had high intensity modifications.

Small cores predominate at VP (85%): more than 58% are bipolar, of which 15 out of 17 (88%) are depleted cores and can no longer yield useable flakes. Bipolar reduction was not detected in the bigger cores. Outside VP, big cores predominate (51%), followed by medium-sized (31%) and small (18%) cores. The percentage of bipolarity increases as the size of cores decreases: 5% in big ones, 62% in medium-sized ones and 71% in small ones. If both sectors are considered together, the minimum number of flake scars varies between two and four, with an average of 3.2 per core. As for the condition of the cores, the large majority (82%) present high intensity modifications, followed by those with medium intensity modification (11%) and only one with minor intensity modification.

### 3.4. Temporal trends

The sites of the Pampean coast analysed in this paper are located in blowout depressions on the dune barrier, where no organic material such as bone or charcoal is preserved and anthropic shellfish accumulations (shell middens) have not been generated; therefore, no absolute radiocarbon dates are available. However, on the basis of Holocene sea level global variations and the absence or presence of several diagnostic features (*e.g.*, pottery and small triangular projectile points) the materials of the dune barrier were assigned to the Middle and late Holocene (Bonomo 2005: 249). In the three assemblages the frequency of bipolar cores is high and even (between 84% and 93%).

On the Patagonian northcentral coast, where the three temporal blocks are represented, the frequency of bipolar cores varies in time (Table 3). For the Middle Holocene (7000 to 2000 years BP), bipolarity appears in low proportion (6%); from 2000 BP on, it increases significantly and progressively, reaching a 54% by the final late Holocene (Table 3).

Table 3. Patagonia northcentral coast: frequency of bipolar cores in temporal block

Temporal Blocks	N= cores	N= bipolar cores	Frequency bipolarity
Middle Holocene	47	3	6,3%
Late Holocene	54	26	48,1%
Final late Holocene	11	6	54,5%

#### 4. Discussion and integration of the data

The selection criteria and technological strategies chosen by hunter-gatherer groups at the analyzed coastal areas can be discussed through the comparison of the local supply of raw materials and the presence of bipolar cores. The results show that bipolar flaking was the main reduction technique applied at the Pampean coast and was very frequent in northcentral Patagonia. Certain differences were noted between the areas. On the Pampean coast, the morphological diversity is low and almost every core is bipolar (87%), even in rocks of medium flaking quality. In northcentral Patagonia, the core type diversity is greater, with bipolar cores representing 50% of the total at the Valdés Peninsula and 39% outside it. As for pebble dimensions, the average size in the Pampean sector is higher: 9 cm as opposed to 7cm for the northcentral Patagonian coast. According to Borrero and Franco, the minimum size that pebbles must have to enable bipolar flaking is 3 cm (Franco & Borrero 1999), so the lithic raw materials from the areas studied would not have prevented a successful application of the bipolar technique. Another possibility could be related to the shape of the nodules. On the Pampean coast, most of the pebbles and cobbles lack natural flat surfaces, which makes them unsuitable for reduction by direct freehand percussion. On the contrary, on the northcentral Patagonian coast there is a greater supply of flattened and faceted gravels, with some flat faces, which could have been used as striking platforms and favoured direct freehand percussion. On the northcentral Patagonian coast the bipolar technique was almost exclusively applied to small and medium-sized pebbles and cobbles from rocks of very good to excellent quality, such as flint, basalts, chalcedonies, fossilized woods and obsidians. Bipolar flaking was not observed in small or medium-sized pebbles of medium quality, such as rhyolites. This would therefore corroborate the hypothesis as concerns size, but the data presented and discussed here shows that the shape of rounded nodules was also taken into account. The differential quality of the raw materials seems to have been considered only on the northcentral coast.

As for temporal trends, because the artefact assemblages of the Pampean coast discussed here do not present absolute chronologies, information from stratified archaeological sites located nearby was taken into account. One of these contexts is the Alfar site, located on the right bank of the Corrientes stream and dated on 5700 <sup>14</sup>C years BP. From a total 251 cores, 98.8% are coastal pebbles and cobbles reduced by the bipolar technique. Basalt prevails (93%), among which 90% present good or very good flaking quality (Apolinaire & Silva 2012). Another adjacent archaeological site is Nutria Mansa 1, which yielded radiocarbon dates between 2705 and 3080 <sup>14</sup>C years BP (Bonomo 2005: 196). It was determined that 92% of the 48 cores recovered are bipolar. The great majority come from coastal pebbles and cobbles, among which basalts (40%) and fine-grained quartzites (22%) prevail. As regards later contexts, at the Claromecó 1 archaeological site, dated on 800 <sup>14</sup>C years BP (Bonomo *et al.* 2008), 100% of the 38 cores are bipolar cores made of coastal pebbles, mainly of basalt and flint of good to very good quality. These data indicate that on the Pampean coast the bipolar flaking of beach pebbles and cobbles - mainly basalts - was a very frequent practice and was sustained over millennia, and increased towards the late Holocene.

For the Pampean coast it has been proposed cultural changes towards the late Holocene such as a process of diversification and intensification in the exploitation of fauna, demographic growth, less frequency trips to the sources, the appearance of technological innovations, and the development of wide networks of social interaction and exchange among human groups from different geographical areas (Berón 2007; Bonomo 2005: 295; González 2005; Martínez & Gutiérrez 2004; Mazzanti 2006; Politis *et al.* 2001). This increasing in the exploitation of the local coastal pebbles (a second class lithic resource in the region) by the bipolar technique, can represent a technological correlate of this process of diversification and intensification, showing less frequent trips to the distant Sierras Bayas quarries, where better quality quartzites were available (Bonomo *et al.* 2008). This increase was also observed in the northcentral Patagonian coast for the same period. In this respect, Gómez Otero (2006) hypothesized that the increase of bipolar core technology towards the final late Holocene could be related with the need to optimize the use of raw materials in a context of demographic growth, reduced mobility and territorial constriction. Therefore, this temporal change in lithic core technology in both areas seems to respond to social matters, technological choices, mobility reduction and new territorial organizations rather than technological needs linked to variations in the availability of coastal raw material.

Temporal and spatial variations can be observed in the application of bipolar flaking in other sectors of the continental Atlantic coastline (Figure 1). In southwest Buenos Aires, province, Bayón and her team detected an almost exclusive application of this technique on local basalt nodules at Middle Holocene contexts, as 58 of the 62 (94%) of the cores recovered from the Las Dunas site are bipolar (Bayón *et al.* 2012). At the mouth of the Colorado river, the use of small (between 2 to 10 cm) local pebbles of flint and basalt was documented, with bipolar flaking being more frequent in the initial Late Holocene contexts (Santos Valero & Armentano 2017). As an example, at the La Primavera site, 67% (n=12) of the bipolar cores are flint and basalt pebbles, followed by quartzites (Bayón *et al.* 2004). At the Las Olas site in the San Blas area, Eugenio & Aldazábal (2004) mention that the cores are mainly bipolar (86.2%) and made from basalt pebbles smaller than 10 cm. On the Río Negro coast of San Matías Gulf, Alberti (2013) documented the use of small bipolar cores throughout the occupational sequence -from the Middle Holocene until the recent temporal block- although in low frequencies (13%) and decreasing later (9%). In the San Jorge gulf, bipolar flaking has not been recorded (Borrero & Caviglia 1978; Arrigoni *et al.* 2008), while only one out of 67 cores analyzed in late Holocene sites of the Santa Cruz province is bipolar (1.5%) and made from non-local raw material: black obsidian from the PDAI source (Ambrústolo 2013: 82). In southern Patagonia there is no record of bipolar cores at Monte León (Cañete Mastrángelo 2016; Cruz & Caracotche 2008), and their frequency is low in Punta Bustamante (15%) where it is applied to small local fossilized wood stones (Mansur 2008).

On the basis of bipolar core frequencies recorded by different researchers, four ranges were established: high (100 to 60%); medium (59 to 30%); low (29 to 10%) and very low to null (9 to 0%). The sectors with the highest values are the coast of the *Interserrana* area between Cape Corrientes and the mouth of the Quequén Salado river; San Blas Peninsula, the mouth of the Colorado and Negro rivers and the northcentral coast of Patagonia (between 60 and 100%). Medium to low frequencies (59 to 10%) are found in the Río Negro coast of the San Matías gulf, and very low to null (<9%) on the coastline that includes the southern Nuevo gulf, almost all of the San Jorge gulf and the area between Monte León and Punta Bustamante in southern Patagonia.

## 5. Synthesis and conclusions

The results of the technological analysis performed on the cores from two different and distant areas of the continental Atlantic coast determined that the most influential variables for the application of bipolar flaking were the small size of nodules and their round shape, generally thick and with no flat natural surfaces. As regards raw material quality, the Pampean coast displayed bipolar flaking in rocks of very good to medium quality, while on the northcentral coast of Patagonia it was applied only in rocks of the best flaking quality. Furthermore, a progressive increase was documented in the region from the Middle Holocene (6%) until the final late Holocene (>54%), that is likely associated with the occurrence of social and organizational changes.

The record of bipolar flaking varies among the different sectors along the Pampa-Patagonia region. In the northern sectors it was applied on small basalt pebbles. On the northcentral Patagonian coast bipolar flaking used on small pebbles of both basalt and flint, and further south (northern coast of Santa Cruz and Punta Bustamante) on pebbles of very high to excellent quality, some non-local (such as obsidian) and some of local but limited availability: such as fossilized woods and dark fine-grained rocks. Therefore, this evidence allows us to infer that bipolar flaking or reduction was a technical strategy applied by hunter-gatherer groups of the Pampa-Patagonia region almost exclusively at those areas where raw material was mainly available in the shape of predominantly small pebbles. At those places where availability was more diversified (with supply at primary and secondary sources), bipolar technique was applied to a lesser extent and restricted to local rocks of the best quality. Finally, the differences observed in the frequency of bipolarity along the Pampa-Patagonia maritime coast suggest it was linked with technological choices that different coastal populations made for the best use of small-sized pebbles through the bipolar technique.

## Acknowledgements

Financial support was provided by diverse grants from the National Research Council of Argentina (CONICET): PIP 0321, 02786, 6470, and 11401000100210 and the Grant 5453/95 from the National Geographic Society granted to Julieta Gómez Otero, and the project “*Arqueología de ambientes acuáticos del Centro-este argentino*” (UNLP Code 11/N885), directed Mariano Bonomo. The authors wish to thank Otis Crandell and two anonymous reviewers for their useful comments, which improved the manuscript. The authors also thank Ana Castelli for corrections to English, and Liliana Calfuquir for bibliography review.

## References

- Alberti, J. 2013, Materias primas líticas y manufactura de instrumentos en tres localidades de la costa oeste del golfo San Matías (provincia de Río Negro, Argentina). *Revista Chilena de Antropología*, 28: 79-105. (in Spanish) (“Lithic Raw Materials and Artifact Manufacture at Three Archaeological Localities in the Western Coast of San Matías Gulf, Río Negro Province, Argentina”) DOI: <https://doi.org/10.5354/0719-1472.2013.30619>
- Ambrústolo, P. 2013, *Estudio de las estrategias de aprovisionamiento y utilización de los recursos líticos por grupos cazadores recolectores en la Costa Norte de Santa Cruz (Patagonia Argentina)*. Unpublished Doctoral thesis at the Facultad de Ciencias Naturales y Museo, Universidad Nacional de La Plata, La Plata, 407 p. (in Spanish)

- (“Study of the supply and use strategies of lithic resources by hunter-gatherer groups on the North Coast of Santa Cruz (Argentine Patagonia)”)
- Ameghino, F. 1910, Une nouvelle industrie lithique: L’industrie de la pierre fendue dans le tertiaire de la région littorale au sud de Mar del Plata. *Anales del Museo Nacional de Buenos Aires* 20(13-3): 189-204. (in French) (“A new lithic industry: The split stone industry in the tertiary of the coastal region south of Mar del Plata”)
- Apolinaire, E. & Silva, C. 2012, Estudios de tecnología lítica en el sitio arqueológico Alfar (partido de General Pueyrredón, provincia de Buenos Aires). *Intersecciones en Antropología*, 13: 525-536. (in Spanish) (“Lithic Technology Studies at the Alfar Archaeological site, General Pueyrredón, Buenos Aires Province”)
- Arrigoni, G., Andrieu, M. & Bañados, C. 2008, Arqueología de cazadores-recolectores prehistóricos en la costa central del golfo San Jorge. In: *Arqueología de la costa patagónica. Perspectivas para la conservación, Río Gallegos* (Cruz, I. & Caracotche, M.S., Eds.), Universidad Nacional de la Patagonia Austral, Subsecretaría de Cultura de la Provincia de Santa Cruz, Santa Cruz: p. 91-107. (in Spanish) (“Prehistoric hunter-gatherer archaeology on the central coast of the San Jorge Gulf”)
- Arthur, K. 2010, Feminine Knowledge and Skill Reconsidered: Women and Flaked Stone Tools. *American Anthropologist*, 112(2): 228-243. DOI: <https://doi.org/10.1111/j.1548-1433.2010.01222.x>
- Aschero, C.A. 1975, *Ensayo para una Clasificación Morfológica de Artefactos Líticos aplicada a Estudios Tipológicos Comparativos*. Informe al Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Buenos Aires, 62 p. (in Spanish) (“Essay for a Morphological Classification of the Lithic Artifacts Applied to Comparative Typological Studies”)
- Aschero, C.A. 1983, *Ensayo para una Clasificación Morfológica de Artefactos Líticos aplicada a Estudios Tipológicos Comparativos. Apéndice A-C. Revisión 1983*. Cátedra de Ergología y Tecnología, Facultad de Filosofía y Letras, Universidad de Buenos Aires, Buenos Aires, 39 p. (in Spanish) (“Essay for a Morphological Classification of the Lithic Artifacts Applied to Comparative Typological Studies. Appendix A-C”)
- Banegas, A. 2016, *La organización de la tecnología lítica y el uso del espacio en la costa centro-septentrional de Patagonia*. Doctoral thesis at the Facultad de Ciencias Naturales y Museo, Universidad Nacional de La Plata, La Plata, 294 p. (in Spanish) (“The organization of lithic technology and the use of space on the central-northern coast of Patagonia”)
- Banegas, A., Goye, S. & Gómez Otero, J. 2016, Caracterización regional de recursos líticos en el nordeste de la provincia del Chubut (Argentina). *Intersecciones en Antropología*, 16(1): 39-50. (in Spanish) (“Regional characterization of lithic resources in the northeast of the province of Chubut (Argentina)”)
- Bayón, C. & Zavala, C. 1997, Coastal sites in south Buenos Aires: a review of Piedras Quebradas. *Quaternary of South America and Antarctic Peninsula*, 10: 229-253.
- Bayón, C., Martínez, G.A., Armentano, G. & Scabuzzo, C. 2004, Arqueología del valle inferior del río Colorado. El sitio La Primavera. *Intersecciones en Antropología*, 5: 39-53. (in Spanish) (“Archaeology of the Lower Valley of the Colorado River. La Primavera site”)



- Bayón, C., Frontini, R. & Vecchi, R. 2012, Middle Holocene settlements on coastal dunes from Southwest of Buenos Aires province, Argentina. *Quaternary International*, 256: 54-61. DOI: <https://doi.org/10.1016/j.quaint.2011.07.014>
- Berón, M. 2007, Circulación de bienes como indicador de interacción entre las poblaciones de la Pampa Occidental y sus vecinos. In: *Arqueología en las Pampas* (Bayón, C., Pupio, A., González, M.I., Flegenheimer, N. & Frére, M., Eds.), Sociedad Argentina de Antropología, Buenos Aires: p. 345-364. (in Spanish) (“Circulation of goods as an indicator of interaction between the populations of the Western Pampa and its neighbours”)
- Berón, M., Migale, L. & Curtoni, P. 1995, Hacia una definición de una Base Regional de Recursos Líticos en el área del Curacó. Una cantera taller: Puesto Córdoba (La Pampa, Argentina). *Relaciones de la Sociedad Argentina de Antropología* 20: 111-128. (in Spanish) (“Towards a definition of a Regional Base of Lithic Resources in the Curacó area. A workshop quarry, La Pampa, Argentina”)
- Binford, L. & Quimby, G. 1963, Indian Sites and Chipped Stone Materials in the Northern Lake Michigan Area. *Fieldiana. Anthropology*, 36: 277-307.  
URL: <https://www.jstor.org/stable/29782305>
- Bonomo, M. 2002, El Hombre Fósil de Miramar. *Intersecciones en Antropología*, 3: 69-85. (in Spanish) (“The Fossil Man from Miramar”)
- Bonomo, M. 2005, *Costeando las Llanuras. Arqueología del Litoral Marítimo Pampeano*. Sociedad Argentina de Antropología, Buenos Aires, 334 p. (in Spanish) (“Coasting the Plains. Archaeology of the Pampean Marine Litoral”)
- Bonomo, M. 2006, Un acercamiento a la dimensión simbólica de la cultura material en la región pampeana. *Relaciones de la Sociedad Argentina de Antropología*, 31: 89-115. (in Spanish) (“An approach to the symbolic dimension of material culture in the Pampas region”)
- Bonomo, M. & Leon, C. 2010, Un contexto arqueológico en posición estratigráfica en los médanos litorales. El sitio Alfar (Pdo. Gral. Pueyrredón, Pcia. Bs. As). In: *Mamül Mapu pasado y presente desde la arqueología pampeana* (Berón, M., Luna, L., Bonomo, M., Montalvo, C., Aranda, C. & Carrera Aizpitarte, M., Eds.), Editorial Libros del Espinillo, Ayacucho: p. 29-45. (in Spanish) (“An archaeological context in a stratigraphic position in the coastal dunes. The Alfar site, Pdo. Gral. Pueyrredón, Pov. Bs. As.”)
- Bonomo, M. & Prates, L. 2014, La explotación de depósitos secundarios de rodados en el curso medio del río negro y el litoral marítimo pampeano. In: *Lithic Artefacts, Mobility and Site Functionality: Problems and Perspectives* (Escola, P.S. & Hocsman, S., Eds.), BAR International Series Vol. 2628, Archaeopress, Oxford: p. 77-92. (in Spanish) (“The exploitation of secondary deposits of rocks in the middle course of the Rio Negro and the Pampa maritime coast”)
- Bonomo, M., Leon, C., Turnes, L. & Apolinaire, E. 2008, Nuevas investigaciones sobre la ocupación prehispánica de la costa pampeana en el Holoceno tardío: el sitio arqueológico Claromecó 1 (Pdo. de Tres Arroyos, Pcia. de Bs. As.). *Intersecciones en Antropología*, 9: 25-41. (in Spanish) (“New investigations on the pre-Hispanic occupation of the Pampas coast in the late Holocene: the Claromecó 1 archaeological site, Pdo. Tres Arroyos, Buenos Aires”)

- Bonomo, M., Scabuzzo, C. & Leon, C. 2013, Cronología y dieta en la costa atlántica pampeana, Argentina. *Intersecciones en Antropología*, 14: 5-18. (in Spanish) ("Chronology and diet on the Atlantic coast of Pampas, Argentina")
- Bórmida, M. 1964, Arqueología de la costa Nordpatagónica. *Trabajos de Prehistoria* 14: 1-108. (in Spanish) ("Archaeology of the Nordpatagónica coast")
- Borrero, L. 1995, Arqueología y paleoecología en Ultima Esperanza: notas para su integración regional. *Cuadernos del Instituto Nacional de Antropología y Pensamiento Latinoamericano* 1994: 185-198. (in Spanish) ("Archaeology and paleoecology in Ultima Esperanza: notes for its regional integration")
- Borrero, L. & Caviglia, S. 1978, *Estratigrafía de los concheros de Bahía Solano: campaña 1976-77*. Actas del libro de resumen del V Congreso Nacional de Arqueología Argentina, San Juan. 1 p. (in Spanish) ("Stratigraphy of the Shell middens of Bahía Solano: 1976-77")
- Cañete Mastrángelo, D. 2016, La Tecnología lítica de la desembocadura del Río Santa Cruz. Una aproximación desde el análisis de los conjuntos artefactuales de los Puntos 37 y 96 de Punta Entrada. *Arqueología*, 22: 311-333. (in Spanish) ("The lithic technology of the mouth of the Santa Cruz River. An approach from the analysis of the artifactual assemblies of Points 37 and 96 of Punta Entrada")
- Codignotto, J., Kokot, R. & Marcomini, S. 1992, Neotectonism and sea-level changes in the coastal zone of Argentina. *Journal of Coastal Research*, 8: 125-133.
- Conlazo, D. 1983, Resultados de una prospección en la zona medanosa de la costa sur de la Provincia de Buenos Aires. In: *Asociación de Estudios Histórico-Arqueológicos de la Región Pampeana*, 2: 32-51, (in Spanish) ("Results of a survey in the medanosa zone of the southern coast of the Province of Buenos Aires")
- Cruz, I. & Caracotche, S. (Eds.) 2008, *Arqueología de la costa patagónica. Perspectivas para la conservación*. Universidad Nacional de la Patagonia Austral-Subsecretaría de Cultura de la Provincia de Santa Cruz, Río Gallegos, 296 p. (in Spanish) ("Archaeology of the Patagonian coast. Conservation prospects")
- Curtoni, R. 1996, Experimentando con bipolares: Indicadores e Implicancias arqueológicas. *Relaciones*, 21: 187-214. (in Spanish) ("Experimenting with bipolar: Indicators and archaeological implications")
- De Aparicio, F. 1932, Contribución al estudio de la arqueología del litoral atlántico de la provincia de Buenos Aires. *Boletín de la Academia Nacional de Ciencias*, 32: 1-180. (in Spanish) ("Contribution to the study of the archaeology of the Atlantic coast of the province of Buenos Aires")
- Díaz de Chiri, M. 1977, Informe preliminar sobre el yacimiento arqueológico Cueva del Tigr (Partido de Necochea, Provincia de Buenos Aires). *Actas y Memorias del IV Congreso Nacional de Arqueología Argentina*, 2: 59-68. (in Spanish) ("Preliminary report on the Cueva del Tigre archaeological site (Necochea, Province of Buenos Aires)")
- Eugenio, E. & Aldazabal, V. 2004, Los cazadores recolectores del litoral marítimo del área de Bahía de San Blas, provincia de Buenos Aires. In: *Contra viento y marea. Arqueología de Patagonia* (Civalero, M., Fernández, P. & Guráieb, G., Eds.), Instituto Nacional de Antropología y Pensamiento Latinoamericano, Buenos Aires: p. 687-700. (in Spanish) ("Hunter-gatherers of the coastline of the Bay area of San Blas, province of Buenos Aires")

- Flegenheimer, N., Bayon, C. & González de Bonaveri, M. 1995, Técnica simple, comportamientos complejos: la talla bipolar en la arqueología bonaerense. *Relaciones de la Sociedad Argentina de Antropología* 20: 81-110. (in Spanish) (“Simple technique, complex behaviors: bipolar carving in Buenos Aires archaeology”)
- Flenniken 1981, *Replicative systems analysis: A model applied to the vein quartz artifacts from the Hoko Riversite*. Laboratory of Anthropology reports 59. Washington State University, Laboratory of Anthropology, Washington, 129 p.
- Flood, J. 1980, *The Moth Hunters. Aboriginal Prehistory of the Australians Alps*. Australian Institute of Aboriginal Studies, Canberra, 388 p.
- Franco, N. & Aragón, E. 2004, Variabilidad en fuentes secundarias de aprovisionamiento lítico: el caso del sur del lago argentino (Santa Cruz, Argentina). *Estudios Atacameños*, 28: 71-85. (in Spanish) (“Variability in secondary lithic supply sources: the case of southern Argentine lake”)
- Franco, N. & Borrero, L. 1999, Metodología de análisis de la estructura regional de recursos líticos. In: *Los tres reinos: Prácticas de recolección en el cono Sur de América* (Aschero C., Korstanje, A. & Vuoto, M., Eds.), Magna Publicaciones, San Miguel de Tucumán: p. 27-37. (in Spanish) (“Methodology for analyzing the regional structure of lithic resources”)
- Gómez Otero, J. 1996, Bases para una arqueología de la costa patagónica central. *Arqueología*, 5: 61-103. (in Spanish) (“Bases for an archaeology of the central Patagonian coast”)
- Gómez Otero, J. 2006, *Recursos, dieta y movilidad en la costa centro-septentrional de Patagonia durante el Holoceno medio y tardío*. Doctoral Thesis at the Facultad de Filosofía y Letras, Universidad de Buenos Aires, Buenos Aires, 482 p. (in Spanish) (“Resources, diet and mobility in the central-northern coast of Patagonia during the middle and late Holocene”)
- Gómez Otero, J. 2007, Isótopos estables, dieta y uso del espacio en la costa atlántica centro septentrional y el valle inferior del río Chubut (Patagonia argentina). In: *Arqueología de Fuego-Patagonia. Levantando piedras, desenterrando huesos... y develando arcanos*, (Morello, F., Martinic, M., Prieto, A. & Bahamondes, G., Eds.), Universidad de Magallanes, Punta Arenas: p. 151-161. (in Spanish) (“Stable isotopes, diet and space use in the northern central Atlantic coast and the lower valley of the Chubut river (Argentine Patagonia)”)
- Gómez Otero, J. & Stern, C. 2005, Circulación, intercambio y uso de obsidias en la costa de la provincia del Chubut (Patagonia, Argentina) durante el Holoceno tardío. *Intersecciones en Antropología*, 6: 93-108. (in Spanish) (“Circulation, exchange and use of obsidians on the coast of the Chubut province (Patagonia, Argentina) during the late Holocene”)
- Gómez Otero, J., Weiler, N., Banegas, A. & Moreno, E. 2013, Ocupaciones del Holoceno medio en Bahía Cracker, costa atlántica de Patagonia central. In: *Tendencias teórico-metodológicas y casos de estudio en la arqueología de la Patagonia* (Zangrando, A., Barberena, R., Gil, A., Neme, G., Giardina, M., Luna, M., Otaola, C., Paulides, S., Salgán, L. & Tivoli, A., Eds.), Museo de Historia Natural de San Rafael, San Rafael: p. 77-186. (in Spanish) (“Mid-Holocene occupations in Cracker Bay, Atlantic coast of central Patagonia”)

- Gómez Otero, J., Belardi, J., Súnico, A. & Taylor, R. 1999, Arqueología de cazadores-recolectores en península Valdés (costa central de Patagonia): primeros resultados. In: *Soplando en el viento. Actas de las III Jornadas de Arqueología de la Patagonia* (Belardi, J.B., Fernández, P.M., Goñi, R.A., Guráieb, A.G. & De Nigris, M., Eds.), Universidad Nacional del Comahue, Neuquén: p. 393-417. (in Spanish) (“Hunter-gatherer archaeology in the Valdés peninsula (central coast of Patagonia): first results”)
- González, M. 2005, *Arqueología de alfareros, cazadores y pescadores pampeanos*. Sociedad Argentina de Antropología, Buenos Aires, 368 p. (in Spanish) (“Archaeology of potters, hunters and fishermen of the Pampas”)
- González, M. & Weiler, N. 1994, Argentinian Holocene transgression: Sideral Ages. *Journal of Coastal Research*, 10(3): 621-627.
- Goodyear, A. 1993, Toolkit Entropy and Bipolar Reduction: A Study of Interassemblage Variability among Paleo-Indian Sites in the Northeastern United States. *North American Archaeologist*, 14: 1-23.
- Guyodo, J. & Marchand, G. 2005, La percussion bipolaire sur enclume dans l’Ouest de la France de la fin du Paléolithique au Chalcolithique: une lecture économique et sociale. *Bulletin de la Société Préhistorique Française*, 102(3): 539-550. (in French) (“TRANSLATION”) DOI: <https://doi.org/10.3406/bspf.2005.13141>
- Haller, M. 1981, *Descripción Geológica de la Hoja 43h “Puerto Madryn”, Provincia del Chubut*. Boletín 184, Servicio Geológico Nacional, Buenos Aires, 41 p. (in Spanish) (“Geological Description of Sheet 43h, Puerto Madryn”)
- Hayden, B. 1980, Confusion in the Bipolar World: Bashed Pebbles and Splintered Pieces. *Lithic Technology*, 9(1): 2-7.
- Holmes, W. 1912, Stone Implements of the Argentine littoral. In: *Early Man in South America* (Hrdlička, A., Ed.), Bureau of American Ethnology, Bulletin Vol. 52, Smithsonian Institute, Washington: p. 125-151.
- Hrdlička, A. (Ed.) 1912, *Early man in South America*. Bureau of American Ethnology, Bulletin 52. Smithsonian Institute, Washington, 405 p.
- Inizan, M., Reduron, M., Roche, H. & Tixier, J. 1995, *Technologie de la pierre taillée 4*. Cercle de Recherches et d’Etudes Préhistoriques (CREP), Meudon 199, p.
- Kobayashi, H. 1975, The Experimental Study of Bipolar Flakes. In: *Lithic Technology, Making and Using Stone Tools* (Swanson, E., Ed.), World Anthropology, De Gruyter, Berlin (New York): p. 115-128. DOI: <https://doi.org/10.1515/9783111390376.115>
- Le Brun-Ricalens, F. 2005, Chronique d’une reconnaissance attendue. Outils « carénés », outils « nucléiformes »: nucléus à lamelles. Bilan après un siècle de recherches typologiques, technologiques et tracéologiques. In: *Productions lamellaires attribuées à l’Aurignacien: chaînes opératoires et perspectives technoculturelles. Acte du XIV<sup>e</sup> congrès de l’Union Internationale des Sciences Pré- et Protohistoriques de Liège 2-8 septembre 2001* (Le Brun-Ricalens F. & Bordes J.-G., Eds.), Archéologiques Vol. 1, Musée nationale d’Histoire et d’Art, Luxembourg: p. 23-72.
- Malvicini, L. & Llambías, E. 1974, Geología y génesis del depósito de manganeso Arroyo Verde, provincia del Chubut, República Argentina. In: *V Congreso Geológico Argentino*, Actas Vol. 2, Buenos Aires: p. 185-202. (in Spanish) (“Geology and genesis of the Arroyo Verde manganese deposit, Chubut province, Argentina”)

- Mansur, E. 2008, Arqueología de la zona de Punta Bustamante (Prov. De Santa Cruz, Argentina). In: *Arqueología de la costa patagónica. Perspectivas para la conservación*, (Cruz, I. & Caracotche, S., Eds.), Universidad Nacional de la Patagonia Austral-Subsecretaría de Cultura de la Provincia de Santa Cruz, Río Gallegos: p. 173-193. (in Spanish) ("Archaeology of the Punta Bustamante area, Prov. de Santa Cruz, Argentina")
- Martínez, G. & Gutiérrez, M. 2004, Tendencias en la Explotación Humana de la Fauna durante el Pleistoceno Final-Holoceno en la Región Pampeana (Argentina). In: *Zooarchaeology of South America* (Mengoni Goñalons, G.L., Ed.), BAR International Series Vol. 1298, BAR Publishing, Oxford: p. 81-98. (in Spanish) ("Trends in Human Exploitation of Fauna during the Final Pleistocene-Holocene in the Pampean Region (Argentina)")
- Mazzanti, D.L. 2006, La constitución de territorios sociales durante el Holoceno Tardío. El caso de las sierras orientales de Tandilia, Argentina. *Relaciones de la Sociedad Argentina de Antropología*, 31: 277-300. (in Spanish) ("The constitution of social territories during the late Holocene. The case of the eastern mountains of Tandilia, Argentina")
- Menghin, O. 1952, Fundamentos cronológicos de la Prehistoria de Patagonia. *Runa*, 5: 23-43. (in Spanish) ("Chronological foundations of the Prehistory of Patagonia")
- Menghín, O. 1963, Industrias de morfología protolítica en Sudamérica. *Anales de la Universidad del Norte*, 2: 69-77. (in Spanish) ("Protolytic morphology industries in South America")
- Mourre, V., & Jarry, M. 2011, « *Entre le marteau et l'enclume... » La percussion directe au percuteur dur et la diversité de ses modalités d'application. Actes de la table ronde organisée à Toulouse du au mars. La Société des Amis du Musée National de Préhistoire et de la Recherche Archéologique avec le concours de l'Université de Toulouse le Mirail. PALEO, numéro spécial 2009-2010. PUBLISHER, Les Eyzies de Tayac, 293 p.* (in French) ("Between the hammer and the anvil..." Direct percussion with a hard striker and the diversity of its methods of application")  
DOI: <https://doi.org/10.4000/paleo.1873>
- Nami, H. 1992, El subsistema tecnológico de la confección de instrumentos líticos y la explotación de los recursos del ambiente: una nueva vía de aproximación. *Shincal*, 2: 33-53. (in Spanish) ("Technological subsystem making stone tools and resources exploitation of the environment: a new avenue of approach")
- Nami, H. 2000, Investigaciones actualísticas y piedra tallada. Criterios experimentales para identificar lascas de talla bipolar: su aplicación en la interpretación de artefactos arqueológicos de los extremos norte y sur de la Patagonia. In: *Actas del III Congreso Argentino de Americanistas 3*, Sociedad Argentina de Americanistas, Buenos Aires: p. 271-292. (in Spanish) ("Current research and carved stone. Experimental criteria to identify flakes of bipolar size: their application in the interpretation of archaeological artifacts from the extreme north and south of Patagonia")
- Orquera, L. & Piana, E.L. 1986, *Normas para la descripción de objetos arqueológicos de piedra tallada. Publicación especial*. Centro Austral de Investigaciones Científicas, Ushuaia, 107 p. (in Spanish) ("Standards for the description of carved stone archaeological objects")



- Orquera, L. & Gómez Otero, J. 2008, Los cazadores-recolectores de las costas de Pampa y Patagonia. *Relaciones de la Sociedad Argentina de Antropología*, 32: 75- 99. (in Spanish) ("The hunter-gatherers of the Pampa and Patagonia coasts")  
URL: <https://dialnet.unirioja.es/descarga/articulo/3100370.pdf>
- Outes, F. 1909, Sobre una Facies Local de los Instrumentos Neolíticos Bonaerenses. *Revista del Museo de La Plata*, 32: 319-339. (in Spanish) ("On a Local Facies of the Buenos Aires Neolithic Instruments")
- Pargeter, J. & Eren, M.I. 2017, Quantifying and Comparing Bipolar Versus Freehand Flake Morphologies, Production Currencies, and Reduction Energetics During Lithic Miniaturization. *Lithic Technology*, 42(2-3): 90-108.  
DOI: <https://doi.org/10.1080/01977261.2017.1345442>
- Politis, G.G. 1984, Arqueología del Área Interserrana Bonaerense. Unpublished Doctoral Thesis at the Universidad Nacional de la Plata, La Plata. (in Spanish) ("Archaeology of the Buenos Aires Interserrana Area")
- Politis, G. & Bayón, C. 1996, Estado actual de las investigaciones en el sitio Monte Hermoso I (provincia de Buenos Aires). *Arqueología*, 6: 83-116. (in Spanish) ("Current status of the investigations at the Monte Hermoso I site (Buenos Aires province)")
- Politis, G., Martínez, G. & Bonomo, M. 2001, Alfarería temprana en sitios de cazadores recolectores de la Región Pampeana (Argentina). *Latin American Antiquity*, 12(2): 167-181. (in Spanish) ("Early pottery at hunter-gatherer sites in the Pampean Region, Argentina")
- Politis, G., Scabuzzo, C. & Tykot, R. 2009, An approach to prehispanic diets in the pampas during early/middle Holocene. *International Journal of Osteoarchaeology* 19: 266- 280.  
DOI: <https://doi.org/10.1002/oa.1053>
- Prous, A.P. 2004, *Apuntes para análisis de industrias líticas*. Ortegalia 2. Monografías de Arqueología, Historia e Patrimonio, Ortigueira, 173 p. (in Spanish) ("Notes for lithic industry analysis")
- Santos Valero, F. & Armentano, G. 2017, La Tecnología Lítica Arqueología de Cazadores-Recolectores del Curso Inferior del río Colorado (Provincia de Buenos Aires, Argentina). In: *Aportes al conocimiento de las ocupaciones humanas Pampeano-Patagónicas* (Martínez, G., Ed.), Series Monográficas INCUAPA-CONICET Vol. 6, CONICEN, Olavarría: p. 123-145. (in Spanish) ("TRANSLATION")
- Shelley, P. 1993, A Geoarchaeological Approach to the Analysis of Secondary Lithic Deposits. *Geoarchaeology: An International Journal*, 8(1): 59-72.  
DOI: <https://doi.org/10.1002/gea.3340080105>
- Shott, J. 1989, Bipolar Industries: Ethnographic Evidence and Archaeological Implications. *North American Archaeologist*, 10: 1-24.
- Sillitoe, P. & Hardy, K. 2003, Living Lithics: Ethnoarchaeology in Highland Papua New Guinea. *Antiquity*, 77: 555-566. DOI: <https://doi.org/10.1017/S0003598X00092619>
- Soriano, S., Robert, A. & Huysecom, É. 2011, Percussion bipolaire sur enclume: choix ou contrainte? L'exemple du Paléolithique d'ounjougou (Pays Dogon, Mali). In: *Entre le marteau et l'enclume... "La percussion directe au percuteur dur et la diversité de ses modalités d'application* (Mourre, V. & Jarry, M., Eds.), PALEO, numéro spécial 2009-2010, Les Eyzies de Tayac: p. 123-132. (in French) ("Bipolar percussion on anvil:

Choice or constraint? The example of the Paleolithic of Ounjougou (Dogon Country, Mali)”)

- Súnico, C. 1996, *Geología del Cuaternario y Ciencia del Suelo: relaciones geomórficas y estratigráficas con suelos y paleosuelos*. Unpublished Doctoral thesis at the Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires, Buenos Aires, 227 p. (in Spanish) (“Quaternary Geology and Soil Science: Geomorphic and Stratigraphic Relationships with Soils and Paleosols”)
- Svoboda, A. 2015, *Los vertebrados pequeños en la subsistencia de cazadores-recolectores: una evaluación zooarqueológica comparativa para Patagonia Central*. Unpublished Doctoral thesis at the Facultad de Filosofía y Letras, Universidad de Buenos Aires, Buenos Aires, 349 p. (in Spanish) (“Small vertebrates in hunter-gatherer subsistence: a comparative zooarchaeological assessment for Central Patagonia”)
- White, P. 1968, Ston naip bilong Tumbuna: The living stone age in New Guinea, In: *La Préhistoire Problèmes et tendances* (Bordes F. & Sonnevile, D., Eds.), Centre national de la recherche scientifique (CNRS), Paris: p. 511-51.

---

# La talla bipolar en la base regional de recursos líticos: Estudio comparativo entre los núcleos de las costas Atlánticas Pampeana y de Patagonia Centro-Septentrional (Argentina)

Anahí Banegas<sup>1</sup>, Mariano Bonomo<sup>2</sup>, Julieta Gómez Otero<sup>1</sup>

1. CONICET - Instituto de Diversidad y Evolución Austral. B. Brown 2915, Puerto Madryn, Argentina.

Email: Banegas: banegas.anahi@yahoo.com.ar; Gómez Otero: julietagomezotero@yahoo.com.ar

2. CONICET - División Arqueología, Facultad de Ciencias Naturales y Museo, Universidad Nacional de La Plata, La Plata, Argentina. Email: mbonomo@fcnym.unlp.edu.ar

---

## Resumen:

La técnica bipolar es una de las primeras técnicas de talla lítica en la historia de la humanidad. Ha sido registrado en sitios arqueológicos de diferentes partes del mundo y de diferentes épocas. Distintas posturas se dieron en relación con las probables causas por las cuales los grupos humanos del pasado lo emplearon, atribuyendo su aplicación a características físicas y naturales de los recursos líticos y a factores sociales relacionados con preferencias estilísticas y tradiciones culturales. Este trabajo presenta los resultados del análisis comparativo de 196 núcleos líticos recuperados en 14 sitios arqueológicos de distinta cronología y sectores de la costa pampeana (entre el Cabo Corrientes y la desembocadura del río Quequén Salado) y la costa centro norte de Patagónica (entre Arroyo Verde y Bahía Cracker). El objetivo de este estudio es evaluar la aplicación de la talla bipolar en ambas áreas y explorar tendencias espaciales y temporales en el uso de esta técnica en otros sectores de la costa atlántica continental desde la provincia de Buenos Aires hasta la provincia de Santa Cruz. Se evaluó el uso de la técnica bipolar en relación con la estructura ambiental lítica, el tamaño de los nódulos y la materia prima. Los núcleos proporcionan información sobre las primeras etapas de la explotación de materias primas líticas para la elaboración de los instrumentos, es decir, la selección inicial y la reducción de los nódulos. La metodología de estudios consistió en muestreos sistemáticos de rocas y el análisis tecnológico de los núcleos. Para el análisis de los núcleos bipolares se registró el tipo de materia prima, el grado de modificación del nódulo original, la longitud, el ancho y el grosor máximos, y la proporción de corteza en cada artefacto. Los resultados obtenidos mostraron altas frecuencias de núcleos bipolares en la costa pampeana (87%) y frecuencias moderadas en la costa norte patagónica central (39%). Se determinó que las variables que más influyeron en la aplicación de la técnica bipolar fueron el pequeño tamaño de los nódulos y su forma redondeada, generalmente espesa y sin superficies planas. En cuanto a la calidad de la materia prima, la técnica se aplicó en rocas de buena a mediana calidad en la costa pampeana, mientras que en la costa norte central se aplicó exclusivamente en rodados de cualidades superiores (muy buenas a excelentes). Se determinó que el registro de la talla bipolar es variable entre los diferentes sectores a lo largo de la costa pampeana y patagónica. En el litoral marítimo de la provincia de Buenos Aires la talla bipolar se aplicó en rodados pequeños de basalto mientras que en el sector de la costa norte de Patagonia central se usó en guijarros de basalto y sílice. En el sector costero de Santa Cruz la talla bipolar se aplicó en guijarros de muy alta a excelente calidad, algunos no locales (como la obsidiana) y en otros de disponibilidad local pero limitada (como xilópalos y rocas oscuras de grano fino). Las diferencias observadas probablemente estén vinculadas con elecciones tecnológicas realizadas por diferentes grupos de cazadores-recolectores para un mejor aprovechamiento de las rocas locales ampliamente disponibles.

**Palabras clave:** técnica bipolar; base de recursos líticos; núcleos; estudio comparativo; costa pampeña y costa atlántica norte y central de Patagonia

### **Extended abstract (English):**

Bipolar reduction is one of the earliest lithic knapping techniques in the history of humankind. It has been recorded in archaeological sites different of parts of the world and from different time periods. Extensive discussions were given in the scientific literature in relation to the reasons why past human groups employed it, attributing its application to physical and natural characteristics of the lithic resources and social factors related to stylistic preferences and cultural traditions. This paper presents the results of the comparative analysis of 196 lithic cores recovered from 14 archaeological sites of diverse chronology along the Pampean coast (between Cape Corrientes and the Quequén Salado river mouth) and the northcentral Patagonian coast (between Verde Stream and Cracker Bay). The aim of this study is to assess the application of bipolar flaking in both areas and to explore spatial and temporal trends of this technique in other sectors of the continental Atlantic coast from Buenos Aires province to Santa Cruz province. The use of the bipolar technique was observed in relation to the lithic environmental structure, the size of nodules and their raw material. To analyse this issues, we also sampled natural deposits of coastal pebbles and cobbles along the Pampas and the Northcentral Patagonia areas. The analysed cores provide information about the first stages of lithic raw material exploitation for tool manufacturing, that is to say, the initial selection and initial reduction of the nodules. For bipolar cores we distinguished the degree of modification suffered by the extremes of the original nodule, in addition to the quality of raw material, maximum length, width and thickness, and the proportion of cortex in each artefact. The obtained results revealed high frequencies of bipolar cores at the Pampean coast (87%) and moderate frequencies at the northcentral Patagonian coast (39%). It was determined that the variables that influenced the application of bipolar techniques the most were the small size of the nodules and their rounded shape, generally thick, and without flat surfaces. As for raw material quality, the technique was applied on good to medium quality rocks at the Pampean coast, while in the northcentral coast it was exclusively applied on pebbles and cobbles of superior qualities (very good to excellent). The record of bipolar flaking varies among the different sectors along the Pampa-Patagonia region. In the Buenos Aires province bipolar flaking was applied on small basalt pebbles and in the northcentral Patagonian coast it was also used on small pebbles of basalt and flint. On the Santa Cruz province it was used on pebbles of very high to excellent quality, some non-local (such as obsidian) and some of local but limited availability (such as fossilized woods and dark fine-grained rocks). The differences observed are most likely linked with technological choices made by different hunter-gatherers to make the best use of widely available local rocks to make tools.